

## [SYSTEMATIC REVIEW]

## OPEN ACCESS

# The role of spirituality, meaning in life, and existential well-being in depression, anxiety, and suicidality: a systematic review

Arveen Kaur<sup>1\*</sup><sup>1</sup> University of Louisville, United States of America\* **Correspondence:** Arveen Kaur, University of Louisville, United States of America. Email: [gorayaarveen93@gmail.com](mailto:gorayaarveen93@gmail.com)

## Abstract

**Background:** Spirituality, meaning in life (MiL), and existential well-being (EWB) have increasingly been investigated as psychological and existential resources relevant to depression, anxiety, and suicidal thoughts and behaviours. However, these constructs are frequently conflated with religiosity, despite evidence that their associations with mental health may differ.

**Objective:** To systematically synthesize empirical evidence concerning associations of spirituality, spiritual well-being, EWB, and MiL with depression, anxiety, suicidal ideation, suicide attempts, and related suicide risk, and to evaluate evidence from interventions explicitly targeting meaning or spiritual/existential well-being.

**Methods:** Searches were conducted on 19 August 2026 across PubMed-indexed literature, Europe PMC, and Semantic Scholar using combinations of terms for spirituality, spiritual well-being, existential well-being, meaning/purpose in life, depression, anxiety, and suicidality. The auditable retrieved screening corpus contained 95 records: 39 PubMed-indexed records, 30 Europe PMC records, and 26 Semantic Scholar records. Three duplicate title/DOI records were removed, 92 records were screened, and 35 primary studies met eligibility criteria. Screening, extraction, and design-specific risk-of-bias appraisal were conducted by one reviewer. Owing to substantial clinical, methodological, and construct heterogeneity, meta-analysis was not considered appropriate and narrative synthesis was performed.

**Results:** Across clinical, community, student, oncology, renal, psychiatric, military/veteran, and older-adult populations, higher EWB and presence of MiL showed the most consistent inverse associations with depression, anxiety, and suicidality. EWB was frequently more strongly associated with mental health than religious well-being. Presence of MiL predicted lower subsequent suicidal ideation in several longitudinal studies, whereas searching for meaning showed heterogeneous and context-dependent associations. General spirituality was usually associated with favourable outcomes but was less consistent after adjustment for social support and other psychosocial factors. Randomized trials of meaning-centered and spiritually oriented interventions demonstrated improvements in meaning/spiritual well-being and selected depressive, anxiety, hopelessness, or desire-for-hastened-death outcomes, although effects were not uniform.

**Conclusions:** Existential meaning, coherence, peace, and purpose appear to constitute clinically relevant protective resources across depression, anxiety, and suicidality. Evidence is strongest for EWB and the presence of MiL rather than for religiosity per se. Nevertheless, causal interpretation is limited by the predominance of observational studies, self-report measurement, cultural heterogeneity, and potential reverse causality. Future longitudinal and mechanism-focused trials should distinguish spiritual, religious, existential, and meaning-related constructs rather than treating them interchangeably.

**Keywords:** spirituality; spiritual well-being; existential well-being; meaning in life; purpose in life; depression; anxiety; suicidal ideation; suicide; systematic review

Received: 22 August 2026

Revised: 27 August 2026

Accepted: 31 August 2026

Published: 01 September 2026

## Cite this article

Kaur A. The role of spirituality, meaning in life, and existential well-being in depression, anxiety, and suicidality: a systematic review. J Evid-Based Med Res. 2026;2(3):31–39. DOI: <https://doi.org/10.66687/jebmr.2.03.2026.38>

© The Author(s) 2026. Published by Acadentium Research. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC BY 4.0), which permits unrestricted use, distribution and reproduction in any medium, provided the original work is properly cited.

## 1. Introduction

Depression, anxiety, and suicidal thoughts and behaviours are multidetermined phenomena arising from interactions among biological vulnerability, psychological processes, interpersonal circumstances, and broader social and cultural factors. Increasing attention has therefore been directed toward potentially protective psychological resources rather than exclusively toward symptoms and risk factors. Spirituality, meaning in life (MiL), purpose, and existential well-being (EWB) are among the constructs most frequently proposed as resources capable of supporting resilience in the face of psychological distress.

These constructs require conceptual separation. Spirituality may involve transcendence, connectedness, inner values, meaning, or relationship with the sacred, whereas religiosity generally refers more specifically to religious beliefs, practices, identities, or institutional participation. Spiritual well-being instruments similarly distinguish religious well-being (RWB), reflecting one's perceived relationship with God or a

transcendent being, from EWB, reflecting purpose, life satisfaction, coherence, and a sense that life is meaningful. MiL research also commonly distinguishes the presence of meaning, indicating that life is already experienced as coherent and purposeful, from the search for meaning, describing active efforts to establish or strengthen that meaning.

Previous systematic reviews have examined religion/spirituality and suicide, existential constructs in suicidal behaviour, or MiL and suicidal ideation separately.[1-3] The latter literature is particularly compelling: a recent meta-analysis of young people reported a substantial inverse association between MiL and suicidal ideation, while also demonstrating considerable methodological diversity among contributing studies.[3] What remains less clearly synthesized is whether the same existential resources relate consistently across the broader continuum of depression, anxiety, and suicidality and whether EWB and meaning differ from explicitly religious dimensions.

The present systematic review therefore aimed to determine: (1) whether spirituality, spiritual well-being, EWB, and MiL are

associated with depression and anxiety; (2) whether these constructs are associated with suicidal ideation, attempts, or suicide risk; (3) whether the presence and search dimensions of MiL show different patterns; (4) whether EWB differs from RWB in its relationship with mental health; and (5) whether interventions targeting meaning or spiritual/existential well-being improve clinically relevant outcomes.

## 2. Methods

### 2.1 Reporting framework

The review was structured according to PRISMA 2020 principles.[4] PRISMA provides standardized guidance for identifying, screening, assessing, and transparently reporting studies in systematic reviews. A prospective protocol was not registered. Because the review was conducted within a single-reviewer research workflow, screening, extraction, and appraisal were performed by one reviewer; the absence of independent duplicate review is acknowledged as a methodological limitation.

### 2.2 Eligibility criteria

Eligibility criteria are summarized in Table 1.

### 2.3 Information sources and search strategy

Searches were performed on 19 August 2026 across three independently searchable literature sources/interfaces: PubMed-indexed literature, Europe PMC, and Semantic Scholar. The core Boolean concept was:

(spirituality OR “spiritual well-being” OR “existential well-being” OR “meaning in life” OR “purpose in life”) AND (depress\* OR anxi\* OR suicid\*)

Syntax was adapted to each interface, with title/abstract-focused terms used where supported. Reference lists of relevant systematic reviews and retrieved primary studies were additionally inspected for potentially relevant papers. Search results were normalized by title and DOI before duplicate removal.

Importantly, the numerical PRISMA counts below represent the auditable records actually retained from the searches run in this review environment.

### 2.4 Study selection

Ninety-five records entered the retrieved screening corpus. Three duplicate title/DOI entries were removed, leaving 92 records for title and abstract screening. Fifty-seven records were excluded because they did not meet the predefined exposure, outcome, design, or primary-research criteria. Thirty-five reports underwent eligibility assessment and were retained in the qualitative synthesis. No report selected for this final eligibility stage could not be retrieved. Complete process is shown in Figure 1.

### 2.5 Data extraction

For each study, data were extracted on country/population, design, relevant spiritual/existential construct, mental-health outcome, principal findings, and key methodological limitations. Particular attention was given to distinguishing EWB from RWB and presence of MiL from search for MiL.

### 2.6 Risk-of-bias assessment

Design-specific domains consistent with established critical appraisal principles were considered: sampling/selection, validity of exposure and outcome measurement, confounder adjustment, temporality, attrition for longitudinal studies, and randomization/blinding/attrition for trials. Because heterogeneous designs make numerical quality scores potentially misleading, studies were classified descriptively as low-to-moderate (L-M), moderate (M), or moderate-to-high (M-H) concern rather than converted into a single numerical score.

## 2.7 Synthesis

Meta-analysis was not performed because studies varied substantially in construct definitions, measurement instruments, populations, study designs, outcomes, and statistical estimates. Results were synthesized narratively by depression/anxiety, suicidality, construct-specific differences, and intervention evidence.

## 3. Results

### 3.1 Study characteristics

The 35 included studies represented diverse populations and research contexts (Table 3). Designs ranged from cross-sectional observational studies to large longitudinal cohorts and randomized psychotherapy/spiritual-care trials. This diversity strengthened the breadth of the evidence but precluded a defensible common pooled effect estimate.

### 3.2 Depression and anxiety

Across medically ill, psychiatric, student, and community populations, EWB was one of the most consistent correlates of lower depression. McCoubrie and Davies found that EWB was inversely associated with both depression and anxiety in advanced cancer whereas RWB was not.[6] Maselko et al. similarly identified a strong inverse relationship between EWB and depression, while religious dimensions displayed materially different associations.[8] Lee’s study of 518 nursing students also found EWB to be more strongly related to lower depression than RWB.[16] These studies argue against treating the religious and existential components of spiritual well-being as interchangeable.

Comparable patterns were observed in advanced cancer and haemodialysis. Spiritual well-being was closely related to psychological distress among advanced cancer patients,[11] while among haemodialysis patients, the FACIT-Sp meaning domain was associated with lower depression, anxiety, and suicide risk, peace with lower depression and anxiety, and faith with selected outcomes; separate measures of religiousness were not associated with the principal mental-health outcomes.[26]

MiL demonstrated a similarly favourable pattern. Steger et al. found that the presence of meaning was associated with lower depression and anxiety,[7] and García-Alandete et al. reported strong negative relationships between MiL and depression, hopelessness, and suicide risk in patients with borderline personality disorder.[15] In newer large studies, MiL was inversely related to adolescent depression and anxiety, with life satisfaction and resilience constituting plausible pathways.[38]

The most recent longitudinal evidence further suggests that directionality may be reciprocal rather than exclusively protective. Tam et al. found longitudinal relationships between MiL and subsequent psychological health among older adults,[35] while Halstead et al. found that presence of meaning predicted lower later depression, anxiety, and self-harm in a large UK cohort, whereas active searching for meaning was associated with some less favourable outcomes.[37] Thus, having meaning and searching for meaning should not be combined as a single exposure.

### 3.3 Suicidal ideation and suicide-related outcomes

Evidence relating MiL and EWB to suicidality was particularly consistent. Among college students, greater spiritual well-being was associated with lower suicidal ideation,[9] while in depressed veterans, life meaning remained related to suicidal ideation even after accounting for depression severity and previous suicidal behaviour.[17] The latter study included 110 depressed veterans and illustrates that meaning may explain variance beyond conventional symptom severity.

Longitudinal data strengthen this observation. Kleiman and Beaver found that meaning predicted changes in suicidal

ideation over follow-up,[13] and studies of older adults showed that MiL predicted lower subsequent suicidal ideation after accounting for depressive symptoms and other risk factors.[21,22] Among Spanish university students, MiL prospectively mediated the association between depressive symptoms and later frequency of suicidal ideation.[34]

The distinction between existential and explicitly religious well-being again emerged in high-risk clinical populations. In African American women who had survived a recent suicide attempt, EWB but not RWB contributed to pathways linking traumatic symptoms with hopelessness and suicidal ideation.[25] Similarly, among patients with bipolar disorder and childhood sexual abuse histories, EWB rather than RWB was identified as the relevant protective dimension.[29]

Population-level findings were more nuanced. Rasic et al. found associations between spirituality/religion and suicidal behaviour in a large representative sample, but some relationships attenuated after accounting for social support.[10] Their subsequent longitudinal epidemiological study likewise found that worship attendance and seeking spiritual comfort did not behave identically across major depression, anxiety disorders, suicidal ideation, and attempts.[12] These results imply that observed spiritual associations can partly reflect interpersonal connectedness and other psychosocial pathways.

### 3.4 Intervention evidence

Three included randomized studies provided evidence that spiritual/existential variables may be modifiable rather than merely correlational markers. Meaning-Centered Group Psychotherapy in advanced cancer improved spiritual well-being and selected depressive, hopelessness, and desire-for-hastened-death outcomes, although anxiety was not consistently improved across analyses.[18] A later individual Meaning-Centered Psychotherapy trial improved meaning and spiritual well-being and demonstrated benefits for selected anxiety and hastened-death outcomes relative to usual care.[27] A spiritually oriented care intervention in haemodialysis patients also reduced depression and anxiety.[33]

These trials support the therapeutic plausibility of meaning and spiritual well-being but do not establish that all spiritual interventions are effective or that meaning itself is the sole mechanism. Psychotherapy trials cannot readily blind participants or therapists, interventions contain multiple active components, and attrition is especially relevant in seriously ill populations. Figure 2 shows this.

### 3.5 Overall evidence pattern

Table 4 summarizes the direction and confidence of the qualitative evidence.

In Figure 3, solid pathways represent the dominant relationships identified in the included evidence. Dashed pathways illustrate modifiers and potential reverse causality.

## 4. Discussion

### 4.1 Principal findings

This systematic review identifies a coherent but qualified pattern: the existential component of spirituality - meaning, purpose, peace, and coherence - shows a more consistent relationship with depression, anxiety, and suicidality than religious identity or religious well-being alone. This conclusion is supported across substantially different populations, including people with cancer, haemodialysis patients, university students, psychiatric patients, military personnel, suicide-attempt survivors, and older adults.

The distinction is clinically important. In several studies that measured both EWB and RWB, only EWB showed an independent protective association.[6,8,25,29] The haemodialysis study by Loureiro et al. similarly demonstrated associations for meaning, peace, and faith subdomains while

separate religiousness measures were not associated with the major outcomes.[26] Consequently, the evidence does not support the simplified assertion that “religion prevents depression or suicide.” Rather, existential coherence, perceived meaning, adaptive spiritual resources, supportive relationships, culture, and religious practices appear to interact.

### 4.2 Why meaning may matter

The included studies suggest several plausible mechanisms. A stable sense of meaning may provide cognitive coherence during adversity, sustain future-oriented goals, increase reasons for living, support hope and resilience, and facilitate adaptive coping. Social connection may contribute separately or mediate some associations between religious/spiritual involvement and psychological health.[10,12] Conversely, depression may reduce motivation, positive future expectancy, and perceived significance of life, thereby eroding meaning. Longitudinal evidence indicating reciprocal associations is therefore theoretically plausible.[35]

The distinction between presence and search for meaning adds further nuance. Presence generally appears beneficial. Search, however, may represent constructive exploration in some contexts but unresolved existential instability in others. Chen et al. demonstrated context-dependent associations according to adverse life events,[32] while more recent longitudinal evidence found search for meaning associated with some poorer mental-health outcomes despite favourable associations for presence.[37] Clinical assessment should consequently avoid interpreting a high search-for-meaning score automatically as psychological strength.

### 4.3 Suicidality

MiL may be particularly relevant to suicidal processes because suicidality involves not only psychiatric symptom severity but also hopelessness, perceived burdensomeness, thwarted future possibilities, and diminished reasons for living. Evidence from veterans, community-dwelling older adults, students, bipolar disorder, and suicide-attempt survivors indicates that meaning-related variables can remain relevant after adjustment for depression or other vulnerabilities.[17,21,22,25,29,34]

These conclusions are compatible with previous systematic and meta-analytic evidence focused specifically on suicide. Prior reviews found heterogeneous but potentially protective associations for religiosity/spirituality,[1] while a 2024 meta-analysis reported a substantial inverse association between MiL and suicidal ideation among young people.[3] The current synthesis extends that conceptual pattern by demonstrating that similar existential resources are associated with depression and anxiety and by emphasizing that EWB and RWB should not be collapsed.

### 4.4 Clinical implications

Spirituality should not be prescribed, imposed, or assumed to be beneficial for every patient. Rather, clinicians may consider assessing whether meaning, purpose, spiritual beliefs, existential concerns, or religious communities are personally important to an individual. For patients who identify these domains as relevant, spiritually integrated or meaning-centered approaches may complement evidence-based psychiatric and psychological care.

Assessment should also consider spiritual struggle rather than only spiritual well-being. Religious guilt, alienation, loss of faith, perceived abandonment, or conflict with a religious community may intensify rather than reduce psychological distress. The mixed findings for explicitly religious dimensions in the included literature reinforce the need for individualized assessment.

### 4.5 Strengths and limitations

The review's principal strengths are its integrative focus across three clinically important outcome domains, explicit distinction

among spirituality, EWB, RWB, presence of meaning, and search for meaning, inclusion of observational and intervention evidence, traceable study-level extraction, and transparent PRISMA flow.

Several limitations require emphasis. First, most included studies were observational, and many were cross-sectional; therefore, associations cannot establish causation. Second, spirituality and meaning were operationalized using heterogeneous instruments, limiting direct comparability. Third, several studies used convenience or specialized clinical samples. Fourth, mental-health outcomes were frequently self-reported. Fifth, cultural and religious contexts varied considerably and may moderate associations. Sixth, publication bias cannot be excluded.

Finally, the search was executed through PubMed-indexed retrieval, Europe PMC, and Semantic Scholar. Native institutional searches of PsycINFO, Embase, Web of Science, and Scopus could not be executed in this environment. The review therefore reports the exact auditable corpus retrieved rather than claiming fictional searches of inaccessible databases. For a journal requiring exhaustive librarian-validated searching of proprietary databases, the same strategy should be rerun through those native interfaces before final submission.

## 5. Conclusions

Spirituality, MiL, and EWB are meaningfully associated with depression, anxiety, and suicidality, but the evidence is most consistent for existential well-being and the presence of meaning in life rather than for religiosity alone. Meaning and existential coherence may help explain why some individuals maintain psychological resilience despite psychiatric, medical, or psychosocial adversity. Longitudinal studies and randomized meaning-centered interventions strengthen, but do not conclusively establish, a causal interpretation.

Future research should distinguish presence from search for meaning, separate EWB from RWB, investigate spiritual struggle and cultural moderation, use repeated longitudinal measurements, and test whether changes in meaning mediate clinical improvement. Spiritual and existential assessment may provide valuable information within person-centred mental-health and suicide-prevention care when incorporated according to the patient's own values and preferences.

## Declarations

### Ethics approval

Not applicable because the study synthesized previously published data.

### Availability of data and materials

The study-level data used for qualitative synthesis are reproduced in the evidence tables of this manuscript.

### Competing interests

None.

### Funding

None.

## References

- Lawrence RE, Oquendo MA, Stanley B. Religion and suicide risk: a systematic review. *Arch Suicide Res.* 2016;20(1):1-21. doi:10.1080/13811118.2015.1004494.
- Costanza A, Prelati M, Pompili M. The meaning in life in suicidal patients: the presence and the search for constructs. A systematic review. *Medicina (Kaunas).* 2019;55(8):465. doi:10.3390/medicina55080465.
- Li J, et al. Meaning in life and suicidal ideation among young people: a systematic review and meta-analysis. *Child Youth Serv Rev.* 2024;158:107477. doi:10.1016/j.childyouth.2024.107477.
- Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ.* 2021;372:n71. doi:10.1136/bmj.n71.

- Coleman CL. The contribution of religious and existential well-being to depression among African American heterosexuals with HIV infection. *Issues Ment Health Nurs.* 2004;25(1):103-110.
- McCoubrie RC, Davies AN. Is there a correlation between spirituality and anxiety and depression in patients with advanced cancer? *Support Care Cancer.* 2006;14(4):379-385. doi:10.1007/s00520-005-0892-6.
- Steger MF, Mann JR, Michels P, Cooper TC. Meaning in life, anxiety, depression, and general health among smoking cessation patients. *J Psychosom Res.* 2009;67(4):353-358. doi:10.1016/j.jpsychores.2009.02.006.
- Maselko J, Gilman SE, Buka S. Religious service attendance and spiritual well-being are differentially associated with risk of major depression. *Psychol Med.* 2009;39(6):1009-1017. doi:10.1017/S0033291708004418.
- Taliaferro LA, Rienzo BA, Pigg RM Jr, Miller MD, Dodd VJ. Spiritual well-being and suicidal ideation among college students. *J Am Coll Health.* 2009;58(1):83-90. doi:10.3200/JACH.58.1.83-90.
- Rasic DT, Belik SL, Elias B, Katz LY, Enns MW, Sareen J. Spirituality, religion and suicidal behavior in a nationally representative sample. *J Affect Disord.* 2009;114(1-3):32-40. doi:10.1016/j.jad.2008.08.007.
- Kandasamy A, Chaturvedi SK, Desai G. Spirituality, distress, depression, anxiety, and quality of life in patients with advanced cancer. *Indian J Cancer.* 2011;48(1):55-59. doi:10.4103/0019-509X.75828.
- Rasic D, Robinson JA, Bolton J, Bienvu OJ, Sareen J. Longitudinal relationships of religious worship attendance and spirituality with major depression, anxiety disorders, and suicidal ideation and attempts: findings from the Baltimore Epidemiologic Catchment Area Study. *J Psychiatr Res.* 2011;45(6):848-854. doi:10.1016/j.jpsychores.2010.11.014.
- Kleiman EM, Beaver JK. A meaningful life is worth living: meaning in life as a suicide resiliency factor. *Psychiatry Res.* 2013;210(3):934-939. doi:10.1016/j.psychres.2013.08.002.
- King M, Llewellyn H, Leurent B, Owen F, Leavey G, Tookman A, et al. Spiritual beliefs near the end of life: a prospective cohort study of people with cancer receiving palliative care. *Psychooncology.* 2013;22(11):2505-2512. doi:10.1002/pon.3313.
- García-Alandete J, Marco Salvador JH, Pérez-Rodríguez S. Predicting role of meaning in life on depression, hopelessness, and suicide risk among Spanish borderline personality disorder patients. *Univ Psychol.* 2014;13(4):1545-1555. doi:10.11144/Javeriana.UPSY13-4.prm.
- Lee Y. The relationship of spiritual well-being and involvement with depression and perceived stress in Korean nursing students. *Glob J Health Sci.* 2014;6(4):169-176. doi:10.5539/gjhs.v6n4p169.
- Braden A, Overholser J, Fisher L, Ridley J. Life meaning is associated with suicidal ideation among depressed veterans. *Death Stud.* 2015;39(1-5):24-29. doi:10.1080/07481187.2013.871604.
- Breitbart W, Rosenfeld B, Pessin H, Applebaum A, Kulikowski J, Lichenthal WG. Meaning-centered group psychotherapy: an effective intervention for improving psychological well-being in patients with advanced cancer. *J Clin Oncol.* 2015;33(7):749-754. doi:10.1200/JCO.2014.57.2198.
- Bamonti P, Lombardi S, Duberstein PR, King DA, Van Orden KA. Spirituality attenuates the association between depression symptom severity and meaning in life. *Aging Ment Health.* 2016;20(5):494-499. doi:10.1080/13607863.2015.1021752.
- Sinclair S, Bryan CJ, Bryan ABO. Meaning in life as a protective factor for the emergence of suicide ideation that leads to suicide attempts among military personnel and veterans with elevated PTSD and depression. *Int J Cogn Ther.* 2016;9(1):87-98. doi:10.1521/ijct.2016.9.1.87.
- Heisel MJ, Flett GL. Does recognition of meaning in life confer resiliency to suicide ideation among community-residing older adults? A longitudinal investigation. *Am J Geriatr Psychiatry.* 2016;24(6):455-466. doi:10.1016/j.jagp.2015.08.007.
- Heisel MJ, Neufeld E, Flett GL. Reasons for living, meaning in life, and suicide ideation: investigating the roles of key positive psychological factors in reducing suicide risk in community-residing older adults. *Aging Ment Health.* 2016;20(2):195-207. doi:10.1080/13607863.2015.1078279.
- Bernard M, Strasser F, Gamondi C, et al. Relationship between spirituality, meaning in life, psychological distress, wish for hastened death, and their influence on quality of life in palliative care patients. *J Pain Symptom Manage.* 2017;54(4):514-522. doi:10.1016/j.jpainsymman.2017.07.019.
- Marco JH, Cañabate M, Pérez S, Llorca G. Associations among meaning in life, body image, psychopathology, and suicide ideation in Spanish participants with eating disorders. *J Clin Psychol.* 2017;73(12):1768-1781. doi:10.1002/jclp.22481.
- Florez IA, Allbaugh LJ, Harris CE, Schwartz AC, Kaslow NJ. Suicidal ideation and hopelessness in African American women: the role of posttraumatic stress and spiritual well-being. *Anxiety Stress Coping.* 2018;31(1):46-58. doi:10.1080/10615806.2017.1369260.
- Loureiro ACT, de Rezende Coelho MC, Coutinho FB, Borges LH, Lucchetti G. The influence of spirituality and religiousness on suicide risk and mental health of patients undergoing hemodialysis. *Compr Psychiatry.* 2018;80:39-45. doi:10.1016/j.comppsych.2017.08.004.
- Breitbart W, Pessin H, Rosenfeld B, Applebaum AJ, Lichenthal WG, Li Y, et al. Individual meaning-centered psychotherapy for the treatment of psychological and existential distress: a randomized controlled trial in patients with advanced cancer. *Cancer.* 2018;124(15):3231-3239. doi:10.1002/cncr.31539.
- Musa AS, Pevalin DJ, Al Khalailah MA. Spiritual well-being, depression, and stress among hemodialysis patients in Jordan. *J Holist Nurs.* 2018;36(4):354-365. doi:10.1177/0898010117736686.
- Lamis DA, Kapoor S, Evans APB. Childhood sexual abuse and suicidal ideation among bipolar patients: existential but not religious well-being as a protective factor. *Suicide Life Threat Behav.* 2019;49(2):401-412. doi:10.1111/sltb.12438.

30. Ibrahim N, Che Din N, Ahmad M, Amit N, Ghazali SE, Wahab S, et al. The role of social support and spiritual wellbeing in predicting suicidal ideation among marginalized adolescents in Malaysia. *BMC Public Health*. 2019;19(Suppl 4):553. doi:10.1186/s12889-019-6861-7.
31. Kassem M, Haddad C, Hallit S, Kazour F. Spirituality, religiosity and suicide risk: associations with mental health outcomes. *Int J Psychiatry Clin Pract*. 2021;25(4):336-343. doi:10.1080/13651501.2020.1787453.
32. Chen Q, Wang XQ, He XX, Ji LJ, Liu MF, Ye BJ. The relationship between search for meaning in life and suicidal ideation in Chinese adolescents: the role of meaning presence and stressful life events. *J Affect Disord*. 2021;282:545-553. doi:10.1016/j.jad.2020.12.156.
33. Durmuş M, Ekinci M. The effect of spiritual care on anxiety and depression in patients receiving hemodialysis treatment: a randomized controlled study. *J Relig Health*. 2022;61(3):2041-2055. doi:10.1007/s10943-021-01386-4.
34. Pérez Rodríguez S, Layrón Folgado JE, Guillén Botella V, Marco Salvador JH. Meaning in life mediates the association between depressive symptoms and future frequency of suicidal ideation in Spanish university students: a longitudinal study. *Suicide Life Threat Behav*. 2024;54(2):286-295. doi:10.1111/sltb.13040.
35. Tam KW, Zhang D, Li Y, Xu Z, Li Q, Zhao Y, et al. Longitudinal relationships between meaning in life and mental health among older adults with multimorbidity. *BMC Geriatr*. 2025;25:195. doi:10.1186/s12877-025-05762-7.
36. Vakili Sadeghi M, Tehrani MAH, Pasha H, Hamidia A, Nabipour M, Khafri S, et al. Spiritual well-being and its association with anxiety and depression among patients with cancer. *PLoS One*. 2025;20(6):e0322923. doi:10.1371/journal.pone.0322923.
37. Halstead I, Heron JE, Haworth CMA, von Gontard A, Joinson CJ. Meaning in life and subsequent mental health and self-harm in a UK birth cohort. *J Affect Disord*. 2026;407:121646. doi:10.1016/j.jad.2026.121646.
38. Jiang L, Zeng X, He H, Wang SB. Meaning in life and adolescent depression and anxiety in China: a large-scale study of dual pathways through life satisfaction and resilience. *BMC Psychol*. 2026;14:150. doi:10.1186/s40359-025-03888-2.
39. Manzouri L, Karami A, Malekzadeh M, Seyed-Nezhad M, Moradi-Joo M. The exploratory relationship between spiritual health and depression, anxiety, and stress in individuals who attempted suicide in Yasuj, Southwest Iran: a cross-sectional study. *BMC Psychiatry*. 2026;26:257. doi:10.1186/s12888-026-07884-7.

## Tables and Figures

**Table 1. Eligibility criteria**

| Domain      | Inclusion criteria                                                                                                                                                             | Exclusion criteria                                                                                                     |
|-------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------|
| Population  | Human participants of any age in community, clinical, student, medical, psychiatric, or other defined populations                                                              | Non-human research                                                                                                     |
| Exposure    | Spirituality, spiritual well-being, EWB, MiL, presence/search for meaning, purpose in life, or explicitly meaning/spirituality-focused intervention                            | Religious affiliation alone without a spirituality/meaning-related construct; unrelated positive-psychology constructs |
| Outcomes    | Depression/depressive symptoms, anxiety/anxiety disorders, suicidal ideation, suicide attempt, suicide risk, desire for hastened death where explicitly related to suicidality | Studies reporting only general quality of life without relevant psychological outcome                                  |
| Designs     | Cross-sectional, case-control, cohort/longitudinal, and randomized or quasi-experimental intervention studies                                                                  | Reviews, editorials, protocols, commentaries, purely theoretical papers                                                |
| Publication | Peer-reviewed, traceable empirical article with sufficient methodological/results information                                                                                  | Abstract-only or non-traceable reports                                                                                 |
| Language    | English-language report available for evaluation                                                                                                                               | Reports without sufficient English information                                                                         |
| Time        | No lower publication-date restriction; searched through 19 August 2026                                                                                                         | Publications after search date                                                                                         |

**Table 2. Search and retrieval audit**

| Source/interface               | Core search concepts                                            | Search date | Records retained in screening corpus |
|--------------------------------|-----------------------------------------------------------------|-------------|--------------------------------------|
| PubMed-indexed literature      | Spirituality/SWB/EBW/MiL/purpose AND depression/anxiety/suicide | 19 Aug 2026 | 39                                   |
| Europe PMC                     | Same concept blocks, syntax adapted                             | 19 Aug 2026 | 30                                   |
| Semantic Scholar               | Same concept blocks, relevance-based retrieval                  | 19 Aug 2026 | 26                                   |
| Total before duplicate removal |                                                                 |             | 95                                   |
| Duplicate title/DOI records    |                                                                 |             | 3                                    |
| Unique records screened        |                                                                 |             | 92                                   |

**Table 3. Characteristics and principal findings of included primary studies**

| Ref.                                                            | Population/design             | Construct                                 | Outcomes                                                     | Principal finding                                                                                       | RoB |
|-----------------------------------------------------------------|-------------------------------|-------------------------------------------|--------------------------------------------------------------|---------------------------------------------------------------------------------------------------------|-----|
| [5] Coleman, 2004; adults with HIV                              | Cross-sectional               | RWB/EBW                                   | Depression                                                   | Spiritual dimensions jointly explained substantial variation in depression                              | M   |
| [6] McCoubrie & Davies, 2006; advanced cancer, n=85             | Cross-sectional               | RWB/EBW                                   | Depression, anxiety                                          | EBW inversely related to both; RWB was not                                                              | M   |
| [7] Steger et al., 2009; smoking-cessation patients             | Cross-sectional               | Presence/search MiL                       | Depression, anxiety                                          | Presence of MiL associated with better psychological health                                             | M   |
| [8] Maselko et al., 2009; community adults, n=918               | Cross-sectional               | EBW/RWB/religious attendance              | Depression                                                   | High EWB strongly associated with lower depression; religious dimensions differed                       | M   |
| [9] Taliaferro et al., 2009; college students, n=457            | Cross-sectional               | Spiritual/EBW                             | Suicidal ideation                                            | Higher spiritual well-being associated with lower ideation                                              | M   |
| [10] Rasic et al., 2009; nationally representative sample       | Cross-sectional               | Spirituality/religion                     | Suicidality                                                  | Several protective associations weakened after social-support adjustment                                | L-M |
| [11] Kandasamy et al., 2011; advanced cancer, n=50              | Cross-sectional               | Spiritual well-being                      | Depression, anxiety/distress                                 | Higher spiritual well-being associated with lower psychological distress                                | M-H |
| [12] Rasic et al., 2011; Baltimore ECA                          | Longitudinal                  | Spirituality/attendance                   | Depression, anxiety, suicidality                             | Prospective relationships varied by religious/spiritual dimension                                       | L-M |
| [13] Kleiman & Beaver, 2013                                     | Longitudinal                  | Presence/search MiL                       | Suicidal ideation/attempt                                    | MiL predicted lower subsequent ideation and lower attempt vulnerability                                 | M   |
| [14] King et al., 2013; palliative cancer                       | Prospective cohort            | Spiritual beliefs                         | Depression, anxiety                                          | Associations were complex and not uniformly protective                                                  | M   |
| [15] García-Alandete et al., 2014; BPD, n=80                    | Cross-sectional               | MiL                                       | Depression, hopelessness, suicide risk                       | MiL strongly inversely related to all three outcomes                                                    | M   |
| [16] Lee, 2014; nursing students, n=518                         | Cross-sectional               | EBW/RWB                                   | Depression/stress                                            | EBW displayed the stronger inverse relationship                                                         | M   |
| [17] Braden et al., 2015; depressed veterans, n=110             | Cross-sectional               | Life meaning                              | Suicidal ideation                                            | Meaning remained associated with ideation after depression/suicide-history adjustment                   | M   |
| [18] Breitbart et al., 2015; advanced cancer, n=253             | RCT                           | Meaning-centered group psychotherapy      | Depression, anxiety, hopelessness, desire for hastened death | Improved SWB and selected depression/hopelessness outcomes; anxiety effect less consistent              | L-M |
| [19] Bamonti et al., 2016; older treatment-seeking adults, n=55 | Cross-sectional moderation    | Spirituality/MiL                          | Depression                                                   | Spirituality attenuated the depression–loss-of-meaning relationship                                     | M-H |
| [20] Sinclair et al., 2016; military/veterans, n=393            | Cross-sectional path analysis | Presence/search MiL                       | SI/attempt                                                   | MiL contributed to pathways separating psychiatric symptoms from suicidality                            | M   |
| [21] Heisel & Flett, 2016; older adults                         | Longitudinal                  | MiL                                       | Suicidal ideation                                            | Baseline MiL conferred prospective resiliency to ideation                                               | L-M |
| [22] Heisel et al., 2016; older adults, 173 baseline            | Longitudinal                  | MiL                                       | Suicidal ideation                                            | MiL explained unique variance beyond depression and loneliness                                          | L-M |
| [23] Bernard et al., 2017; palliative care                      | Cross-sectional               | Spirituality/MiL                          | Distress, wish for hastened death                            | Meaning/spirituality related to distress and hastened-death wishes                                      | M   |
| [24] Marco et al., 2017; eating disorders                       | Cross-sectional               | MiL                                       | Psychopathology/SI                                           | Lower MiL associated with greater psychopathology and SI                                                | M   |
| [25] Florez et al., 2018; suicide-attempt survivors, n=113      | Longitudinal                  | EBW/RWB                                   | Hopelessness/SI                                              | EBW, but not RWB, mediated adverse pathways to SI                                                       | M   |
| [26] Loureiro et al., 2018; haemodialysis                       | Cross-sectional               | Meaning/peace/f aith                      | Depression, anxiety, suicide risk                            | Meaning associated with lower depression, anxiety and suicide risk; religiousness measures were null    | M   |
| [27] Breitbart et al., 2018; advanced cancer, n=321             | RCT                           | Individual meaning-centered psychotherapy | Depression, anxiety, SWB, hastened death                     | Improved meaning and SWB; benefits observed for anxiety and desire for hastened death versus usual care | L-M |
| [28] Musa et al., 2018; haemodialysis                           | Cross-sectional               | SWB/EBW                                   | Depression/distress                                          | Greater spiritual/existential well-being associated with less distress                                  | M   |


| Ref.                                                       | Population/design               | Construct                | Outcomes                       | Principal finding                                                                                     | RoB |
|------------------------------------------------------------|---------------------------------|--------------------------|--------------------------------|-------------------------------------------------------------------------------------------------------|-----|
| [29] Lamis et al., 2019; bipolar disorder, n=112           | Cross-sectional moderation      | EWB/RWB                  | Suicidal ideation              | EWB, but not RWB, acted as a protective factor                                                        | M   |
| [30] Ibrahim et al., 2019; marginalized adolescents, n=176 | Cross-sectional                 | RWB/EWB/SWB              | Suicidal ideation              | Greater SWB and family support predicted lower ideation                                               | M   |
| [31] Kassem et al., 2021                                   | Cross-sectional                 | Spirituality/religiosity | Suicide risk                   | Spirituality rather than religiosity showed the clearer inverse association                           | M   |
| [32] Chen et al., 2021; adolescents, n=1,705               | Cross-sectional/moderation      | Presence/search MiL      | Suicidality                    | Presence was protective; search effects depended on adverse-life-event context                        | M   |
| [33] Durmuş & Ekinci, 2022; n=71                           | Randomized intervention         | Spiritual care           | Depression/anxiety             | Spiritual-care intervention reduced psychological symptoms                                            | M   |
| [34] Pérez Rodríguez et al., 2024; university students     | Longitudinal                    | MiL                      | Depression → future SI         | MiL mediated the association between depression and future SI                                         | M   |
| [35] Tam et al., 2025; older adults, n=1,077               | Longitudinal                    | MiL                      | Depression/anxiety             | Evidence supported reciprocal relationships between MiL and mental health                             | L-M |
| [36] Vakili Sadeghi et al., 2025; cancer, n=200            | Cross-sectional                 | SWB/EWB/RWB              | Depression/anxiety             | Greater spiritual well-being associated with markedly lower symptoms                                  | M   |
| [37] Halstead et al., 2026; UK cohort, n=3,337             | Longitudinal                    | Presence/search MiL      | Depression, anxiety, self-harm | Presence predicted more favourable outcomes; search predicted some poorer outcomes                    | L-M |
| [38] Jiang et al., 2026; Chinese adolescents               | Large cross-sectional mediation | MiL                      | Depression/anxiety             | Higher MiL associated with lower symptoms partly through life satisfaction/resilience                 | M   |
| [39] Manzouri et al., 2026; suicide attempters, n=136      | Cross-sectional                 | Spiritual health         | Depression/anxiety/stress      | Strong inverse associations with depression/stress; association with anxiety was weak/non-significant | M   |

Note. RoB, risk of bias; BPD, borderline personality disorder; EWB, existential well-being; RWB, religious well-being; SWB, spiritual well-being; MiL, meaning in life; SI, suicidal ideation.

**Table 4. Integrated evidence synthesis by construct and outcome**

| Construct                                 | Depression                     | Anxiety                        | Suicidality                                              | Overall interpretation                                     | Principal caveats                                                       |
|-------------------------------------------|--------------------------------|--------------------------------|----------------------------------------------------------|------------------------------------------------------------|-------------------------------------------------------------------------|
| Existential well-being                    | Consistent inverse association | Consistent inverse association | Consistent inverse association                           | Strongest and most reproducible spiritual-domain pattern   | Predominantly self-report; some cross-sectional studies                 |
| General spirituality/spiritual well-being | Generally inverse              | Generally inverse              | Generally inverse                                        | Usually favourable but heterogeneous                       | Construct definitions vary; social support may confound effects         |
| Presence of meaning in life               | Consistent inverse             | Generally inverse              | Consistent inverse/protective longitudinal signal        | Strong evidence across populations                         | Possible bidirectionality                                               |
| Search for meaning                        | Mixed                          | Mixed/occasionally adverse     | Context-dependent                                        | Should not be treated as equivalent to presence of meaning | Search may indicate unresolved distress                                 |
| Religious well-being/religious dimensions | Mixed                          | Mixed                          | Mixed/contextual                                         | Less consistently protective than EWB                      | Culture, attendance, social support, positive/negative religious coping |
| Meaning/spiritual interventions           | Selected improvement           | Selected improvement           | Some improvement in hopelessness/hastened-death outcomes | Promising but not definitive                               | Few RCTs; heterogeneous treatments and outcomes                         |

Figure 1. PRISMA-style flow diagram of study identification and selection.



Counts describe the auditable records retained from searches run on 19 August 2026; they are not claimed as exhaustive hit totals of the underlying bibliographic databases.

The flow diagram reports the auditable screening corpus retrieved on 19 August 2026: 95 records retained, three title/DOL duplicates removed, 92 records screened, 57 excluded, and 35 primary studies included in qualitative synthesis. Counts should not be interpreted as exhaustive global hit counts of the underlying bibliographic databases.

Figure 2. Qualitative evidence-direction matrix for spirituality, existential well-being, meaning in life, and mental-health outcomes.

Qualitative Direction and Consistency of the Included Evidence


|                                             |                     |                     |                     |
|---------------------------------------------|---------------------|---------------------|---------------------|
| Existential well-being                      | Consistent inverse  | Consistent inverse  | Consistent inverse  |
| Overall spirituality / spiritual well-being | Mostly inverse      | Mostly inverse      | Mostly inverse      |
| Presence of meaning                         | Consistent inverse  | Inverse             | Consistent inverse  |
| Search for meaning                          | Mixed / conditional | Mixed / conditional | Mixed / conditional |
| Religious well-being / religious dimension  | Mixed               | Mixed               | Mixed / contextual  |
|                                             | Depression          | Anxiety             | Suicidality         |

Darker cells denote a more consistent inverse association with adverse mental-health outcomes.  
The matrix is a narrative-synthesis summary, not a pooled effect-size heat map.

Darker cells represent greater consistency of inverse associations across included studies. The figure represents narrative synthesis rather than pooled effect sizes.

Figure 3. Conceptual model linking spiritual/existential resources with depression, anxiety, and suicidality.

Conceptual Model Derived from the Narrative Synthesis



Solid arrows indicate the dominant direction supported across the included literature; dashed arrows indicate moderation or plausible reciprocal pathways that limit causal interpretation of cross-sectional findings.

Solid pathways represent the dominant relationships identified in the included evidence. Dashed pathways illustrate modifiers and potential reverse causality.

OPEN ACCESS & COPYRIGHT

© 2026 The Author(s). Published by Acadentium Research on behalf of Journal of Evidence-Based Medical Research. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International (CC BY 4.0) licence, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.  
Publisher's note: Acadentium Research remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.